



**THE OPTIMAL SYMMETRICAL POINTS FOR POLYNOMIAL  
INTERPOLATION OF REAL FUNCTIONS IN THE TETRAHEDRON**

by

**Qi Chen**

and

**Ivo Babuška**

**Technical Note BN-1186**

19951116 026

**July 1995**

**DTIC QUALITY INSPECTED 3**

**Institute for  
Physical Science and Technology**

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Note BN-1186	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) The Optimal Symmetrical Points for Polynomial Interpolation of Rea Functions in the Tetrahedron		5. TYPE OF REPORT & PERIOD COVERED Final Life of Contract
7. AUTHOR(s) Qi Chen <sup>1</sup> and Ivo Babuska <sup>2</sup>		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS <sup>2</sup> Institute for Physical Science and Technology University of Maryland College Park, MD 20742-2431		8. CONTRACT OR GRANT NUMBER(s) <sup>2</sup> ONR/N00014-90-J-1030
11. CONTROLLING OFFICE NAME AND ADDRESS Department of the Navy Office of Naval Research Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE July 1995
		13. NUMBER OF PAGES 10
		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release: distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT  The main result of this paper is the computation of the mean optimal symmetrical interpolation points in the tetrahedron up to degree 9. This interpolation set has the smallest Lebesgue constant known today.		

# The Optimal Symmetrical Points for Polynomial Interpolation of Real Functions in the Tetrahedron

Qi Chen\*

Sachs Freeman Associates  
1401 McCormick Drive  
Landover, MD 20785

Ivo Babuška<sup>†</sup>

Institute of Physical Science and Technology  
University of Maryland  
College Park, MD 20742

August 18, 1995

## Abstract

The main result of this paper is the computation of the mean optimal symmetrical interpolation points in the tetrahedron up to degree 9. This interpolation set has the smallest Lebesgue constant known today.

---

\*Supported by NASA Langley and U.S. Naval Research Laboratory.

<sup>†</sup>Partially supported by the Office of Naval Research under Grant N00014-90-J1030.

High quality polynomial interpolation of functions is essential in many applications [1] [7], in particular, in the p-version of the Finite Element Method. However, very little is known about the approximation accuracy of interpolation in the simplex in several dimensions. Of particular concern are the two dimensional and three dimensional cases. We refer here to the few literatures that discuss this problem [2] [3] [4] [6].

In this paper, we compute the mean optimal interpolation sets in the tetrahedron. The main result in [3] is that in the interval and in the triangle, the mean optimal sets are close to the optimal interpolation sets in the mean norm (the smallest Lebesgue constant sets). We give the mean optimal sets in table 3 in this paper. In the literature, the most widely used interpolation sets in the tetrahedron is the equally spaced point sets. The approximation quality for equally spaced point sets is known to deteriorate considerably for high degree interpolation. Bos [2] has proposed a set for arbitrary dimension up to degree 4. Aside from these, We are unaware of any other high degree interpolation sets in the tetrahedron.

Interpolation in a simplex of dimensions more than three is still of interest even though it is not considered in this paper. The general principle introduced in [3] can still be employed, albeit in a more complicated fashion.

We use the same notation as is used in [3]. Let  $S^3$  be the three dimensional simplex in  $R^3$ . Let  $P_n(S^3)$  be the space of polynomials of degree  $n$  in three variables. Then  $\dim P_n(S^3) = N_n := \binom{n+3}{3} = \frac{(n+1)(n+2)(n+3)}{6}$ . Let  $T^n = (x_1, \dots, x_{N_n})$ ,  $x_j \in S^3$  be a set of distinct points in  $S^3$ , where  $n$  is the degree of interpolation.  $T^n$  will be called the nodal set and  $x_i$  the nodal points. Furthermore let  $\mathcal{T}^n$  be the family of all nodal sets, or the set of admissible nodes (see next section for discussion of admissible nodes) *i.e.*,  $T^n \in \mathcal{T}^n$ . The interpolation problem now reads: Given a continuous function in  $S^3$ ,  $f \in C(S^3)$  and the nodal set  $T^n$ , find  $p_n \in P_n$  such that  $p_n(x_i) = f(x_i)$ ,  $x_i \in T^n(S^3)$ ,  $i = 1, \dots, N_n$ . Let  $m_j(x)$ ,  $j = 1, \dots, N_n$  be the basis functions of  $P_n(S^3)$ , then  $p_n(x_i) = \sum_{j=1}^{N_n} a_j m_j(x_i) = f(x_i)$ ,  $1 \leq i \leq N_n$ . This system

of linear equations is uniquely solvable for all right hand sides if and only if the determinant is nonzero. The determinant of the system is denoted by  $VDM(T^n)$ . Assuming now that  $VDM(T^n) \neq 0$ , we can construct the Lagrange coefficients  $L_j^T(x) \in P_n(S^3)$ ,  $j = 1, \dots, N_n$  as basis of  $P_n(S^3)$  with  $L_j^T(x_i) = \delta_{i,j}$ ,  $1 \leq i, j \leq N_n$ . For  $f \in C(S^3)$ , we have

$$p_n(x) = \sum_{k=1}^{N_n} f(x_k) L_k^T(x). \quad (1)$$

Denote  $p_n(t) = \mathcal{L}_T f$ , then  $\mathcal{L}_T$  is a linear projection operator which maps  $C(S^3)$  onto  $P_n(S^3)$ .

If no misunderstanding will occur, we sometimes omit the index  $n$ .

We equip  $P_n(S^3)$  with a norm  $\|\cdot\|$  and denote

$$\|\mathcal{L}_T\| = \sup_{f \neq 0} \frac{\|\mathcal{L}_T f\|}{\|f\|_\infty}, \quad (2)$$

if  $\|\cdot\| = \|\cdot\|_\infty$ , we write  $\|\mathcal{L}_T\|_\infty = \lambda(T)$ . It is easy to show that:

$$\lambda(T) = \max_{x \in S^3} \sum_{k=1}^{N_n} |L_k^T(x)|. \quad (3)$$

$\lambda(T)$  is called the Lebesgue constant of  $\mathcal{L}_T$ . In addition, we define

$$\|\langle \mathcal{L}_T \rangle\|_2^2 := \int_{S^3} \sum_{k=1}^{N_n} |L_k^T(x)|^2 dx, \quad (4)$$

Now we are interested in the sets  $T_1$  which minimizes  $\|\mathcal{L}_T\|_\infty$  and  $T_2$  which minimizes  $\|\langle \mathcal{L}_T \rangle\|_2$ .  $T_1$  is the optimal interpolation set.  $T_2$  is called the mean  $L^2$  (or briefly mean) optimal set. We are also interested in the set  $T_{VDM}$  which maximizes  $|VDM|$  and call it the VDM set. The VDM set is sometimes used to approximate the optimal set, see [2].

Remark: Let  $f \in C(S^3)$ , then  $\|f - \mathcal{L}_T f\|_\infty \leq (1 + \lambda(T)) \inf_{g \in P_n(S^3)} \|f - g\|_\infty$ , see [3]. Hence, the interpolation error is of the same order as the best approximation up to the Lebesgue constant. Interpolation points with the smallest Lebesgue constant therefore leads to the smallest interpolation error.




Figure 1: The standard tetrahedron.

From now on,  $S^3$  is the standard tetrahedron shown in Fig. 1. We denote  $(x, y, z)$  the Cartesian coordinates.

We shall be interested in the degree  $n$  optimal interpolation set for the mean norm (or the VDM determinant) in a restricted set  $\mathcal{T}$  of the nodal points:

- The set of nodal points is symmetrical with respect to all symmetries of the tetrahedron;
- On each face of the tetrahedron, we use degree  $n$  optimal interpolation set for the mean norm (or the VDM determinant) in the triangle.

Condition b. is based on the following consideration. When we approximate a function in a domain partitioned into tetrahedrons, we require the resulting piecewise polynomial to be continuous. We also want to have minimal error on each face. Condition b. is not a serious impediment. Indeed in the triangle case, we computed the degree  $n$  optimal set  $\tilde{T}_{2Tri}^n$  for the mean norm under the constraint that the nodal points on the edges of the triangle is the one dimensional degree  $n$  mean optimal set. The corresponding Lebesgue constant is quite close to that of the actual mean norm optimal set  $T_{2Tri}^n$ . This is shown in Table 1.

Because finding optimal points  $T_1$  is much more difficult, we minimize  $\|\langle \mathcal{L}_T \rangle\|_2$  and find

Table 1: Lebesgue constant and the mean norm for the triangle optimal mean set  $T_{2Tri}^n$ , and the constrained mean set  $\tilde{T}_{2Tri}^n$ . Note the square of the mean norm is normalized by the triangle area.

$n$	$\lambda(T_{2Tri}^n)$	$\lambda(\tilde{T}_{2Tri}^n)$	$\frac{1}{\sqrt{3}} \ \langle \mathcal{L}_{T_{2Tri}^n} \rangle\ _2^2$	$\frac{1}{\sqrt{3}} \ \langle \mathcal{L}_{\tilde{T}_{2Tri}^n} \rangle\ _2^2$
2	$1\frac{2}{3}$	$1\frac{2}{3}$	$\frac{19}{30}$	$\frac{19}{30}$
3	2.1114	2.1206	0.7404	0.7420
4	2.6920	2.6833	0.8196	0.8220
5	3.3010	3.2695	0.8837	0.8863
6	3.7910	3.7736	0.9408	0.9434
7	4.3908	4.3949	0.9964	0.9990
8	5.0893	5.1106	1.0553	1.0578
9	5.9181	5.9456	1.1227	1.1252
10	7.0850	7.1140	1.2050	1.2076
11	8.3383	8.3864	1.3110	1.3140
12	8.6928	8.7384	1.4533	1.4572
13	12.0464	12.1120	1.6508	1.6561

the optimal set  $T_2$  in the constrained family. More precisely we find a set which is an approximation of a local minimum of  $\|\langle \mathcal{L}_T \rangle\|_2$ . We also compute the maximum of  $|VDM|$  subject to the same constraints. We point out that the tetrahedron VDM set  $T_{VDM}^n$  restricted to a face of the tetrahedron is exactly the two dimensional triangle VDM set computed in [3]. Therefore, constraint b. is automatically satisfied by the VDM set. This can be seen by using the hierarchic shape functions in [7] as the basis functions  $m_j(x)$ . Hierarchic shape functions for the tetrahedron are organized into four categories: the nodal shape functions, the edge modes, the face modes and the interior modes.  $|VDM|$  in the hierarchic shape function basis is the product of  $|VDM|$  for the edge modes,  $|VDM|$  for the face modes and  $|VDM|$  for the interior modes. Therefore, maximizing  $|VDM|$  for the tetrahedron is equivalent to maximizing  $|VDM|$  for each edge, each face and the interior.

By symmetry, a nodal point inside the tetrahedron can be at the center of the tetrahedron, or on one of the four lines passing through a vertex and the center of the triangle face opposite to that vertex, or on one of the three lines connecting the midpoints of two

nonintersecting edges, or on one of the six symmetry planes but on none of the aforementioned lines, or on none of the symmetry planes (this node is located inside one of the twenty four subtetrahedrons bounded by the six symmetry planes). It corresponds to a point in a singlet, or a quartet, or a sextet, or a twelve fold symmetrical point, or a twenty four fold symmetrical point. We denote the number of these multiplets by  $n_1$  ( $n_1 = 0$  or  $1$ ),  $n_4$ ,  $n_6$ ,  $n_{12}$ ,  $n_{24}$ . Since the total number of these multiplets equals the number of interpolating nodes inside the tetrahedron,  $n_1 + 4n_4 + 6n_6 + 12n_{12} + 24n_{24} = \frac{(n-1)(n-2)(n-3)}{6}$ .

The integer solution for the above equation is nonunique when  $n \geq 6$ . Different integer solutions correspond to different symmetry patterns of the nodal set. Each symmetry pattern has a minimum mean norm, we want to find the smallest mean norm (the global minimum) among these minima.

For the  $n$ -th degree equally spaced point set, the symmetry pattern will be denoted as  $(n_1^{eq}, n_4^{eq}, n_6^{eq}, n_{12}^{eq}, n_{24}^{eq})$ . It obviously satisfy the integer equation  $n_1 + 4n_4 + 6n_6 + 12n_{12} + 24n_{24} = \frac{(n-1)(n-2)(n-3)}{6}$ . Numerical evidence shows that the minimum for this symmetry is smaller than the minima in other symmetry pattern cases. Therefore, we restrict ourselves to one more constraint:

c. The integer pair  $(n_1, n_4, n_6, n_{12}, n_{24})$  for the symmetrical pattern of the degree  $n$  optimal set is  $(n_1^{eq}, n_4^{eq}, n_6^{eq}, n_{12}^{eq}, n_{24}^{eq})$ , the solution for the symmetrical pattern of the degree  $n$  equally spaced point set.

In Table 3, we list the approximate optimal sets in the tetrahedron barycentric coordinates:  $b_1 = \frac{1}{2}(1 - x - \frac{y}{\sqrt{3}} - \frac{z}{\sqrt{6}})$ ,  $b_2 = \frac{1}{2}(1 + x - \frac{y}{\sqrt{3}} + \frac{z}{\sqrt{6}})$ ,  $b_3 = \frac{1}{\sqrt{3}}(y - \frac{z}{\sqrt{6}})$ ,  $b_4 = \sqrt{\frac{3}{8}}z$ .  $b_1 + b_2 + b_3 + b_4 = 1$ . We also list  $(n_1^{eq}, n_4^{eq}, n_6^{eq}, n_{12}^{eq}, n_{24}^{eq})$  and the number of interpolation points  $N_n$ . Points with multi-fold symmetry are listed only once. Other points with the same symmetry can be obtained by applying tetrahedron symmetries to the listed points, *i.e.*, by permuting barycentric coordinates  $b_1, b_2, b_3, b_4$ .



Table 2: Lebesgue constant and the mean norm for the optimal mean set, the VDM set and the equally spaced point set. Note the square of the mean norm is normalized by the tetrahedron volume.

$n$	$\lambda(T_2^n)$	$\lambda(T_{VDM}^n)$	$\lambda(T_{eq}^n)$	$\frac{3}{\sqrt{8}}\ \langle \mathcal{L}_{T_2^n} \rangle\ _2^2$	$\frac{3}{\sqrt{8}}\ \langle \mathcal{L}_{T_{VDM}^n} \rangle\ _2^2$	$\frac{3}{\sqrt{8}}\ \langle \mathcal{L}_{T_{eq}^n} \rangle\ _2^2$
2	2	2	2	$\frac{18}{35}$	$\frac{18}{35}$	$\frac{18}{35}$
3	2.9339	2.9329	3.0200	0.6435	0.6437	0.6893
4	4.1120	4.1534	4.8801	0.7669	0.7670	0.9329
5	5.6158	5.9961	8.0937	0.8892	0.8941	1.2915
6	7.3632	8.8898	13.6568	1.0237	1.0423	1.8721
7	9.3659	11.6425	23.3789	1.1895	1.2417	2.9157
8	12.3111	15.8340	40.5455	1.4165	1.5513	4.9854
9	15.6857	22.3304	71.1521	1.7532	2.0984	9.4538

Neither  $T_2^n$  nor  $T_{VDM}^n$  leads to the minimization of the Lebesgue constant. Nevertheless based on the one dimensional and the triangle results, we expect that the Lebesgue constant of  $T_2^n$  will not be significantly larger than the Lebesgue constant of  $T_1^n$ , and will be smaller than the Lebesgue constant of  $T_{VDM}^n$ . In Table 2, we give the Lebesgue constants for the sets  $T_2^n$  and  $T_{VDM}^n$ . For comparison, we also list the Lebesgue constant for the equally spaced point set  $T_{eq}^n = \{(b_1 = \frac{i}{n}, b_2 = \frac{j}{n}, b_3 = \frac{k}{n}, b_4 = \frac{l}{n}), 0 \leq i, j, k, l; i + j + k + l = n\}$ . Indeed,  $T_2^n$  has a smaller Lebesgue constant than  $T_{VDM}^n$  and a much smaller Lebesgue constant than that of  $T_{eq}^n$ . The last three columns give the mean norms.

We have given in this paper the mean optimal set  $T_2^n$  in the tetrahedron.  $T_2^n$  has the smallest Lebesgue constant known today. The other two interpolation sets are the equally spaced point set  $T_{eq}^n$  and the VDM set  $T_{VDM}^n$ .

Table 3: Barycentric coordinates for the tetrahedron optimal mean set  $T_2$ .

$n$	$N_n$	$n_1$	$n_4$	$n_6$	$n_{12}$	$n_{24}$	$b_1$	$b_2$	$b_3$	$b_4$
2	10	0	0	0	0	0	1.0000000	0.0000000	0.0000000	0.0000000
							0.5000000	0.5000000	0.0000000	0.0000000
3	20	0	0	0	0	0	1.0000000	0.0000000	0.0000000	0.0000000
							0.7251957	0.2748043	0.0000000	0.0000000
							0.3333333	0.3333333	0.3333333	0.0000000
4	35	1	0	0	0	0	1.0000000	0.0000000	0.0000000	0.0000000
							0.8306024	0.1693976	0.0000000	0.0000000
							0.5000000	0.5000000	0.0000000	0.0000000
							0.2208880	0.2208880	0.5582239	0.0000000
							0.2500000	0.2500000	0.2500000	0.2500000
5	56	0	1	0	0	0	1.0000000	0.0000000	0.0000000	0.0000000
							0.8866427	0.1133573	0.0000000	0.0000000
							0.6431761	0.3568239	0.0000000	0.0000000
							0.1525171	0.1525171	0.6949657	0.0000000
							0.4168658	0.4168658	0.1662683	0.0000000
							0.1823054	0.1823054	0.1823054	0.4530838
6	84	0	1	1	0	0	1.0000000	0.0000000	0.0000000	0.0000000
							0.9194021	0.0805979	0.0000000	0.0000000
							0.7349105	0.2650895	0.0000000	0.0000000
							0.5000000	0.5000000	0.0000000	0.0000000
							0.3333333	0.3333333	0.3333333	0.0000000
							0.1097139	0.1097139	0.7805723	0.0000000
							0.3157892	0.5586077	0.1256031	0.0000000
							0.1357838	0.1357838	0.1357838	0.5926485
							0.3559336	0.3559336	0.1440664	0.1440664
							0.3559336	0.3559336	0.1440664	0.1440664
7	120	0	2	0	1	0	1.0000000	0.0000000	0.0000000	0.0000000
							0.9398927	0.0601073	0.0000000	0.0000000
							0.7957614	0.2042386	0.0000000	0.0000000
							0.6042138	0.3957862	0.0000000	0.0000000
							0.0817370	0.0817370	0.8365261	0.0000000
							0.4494208	0.4494208	0.1011584	0.0000000
							0.2663399	0.2663399	0.4673202	0.0000000
							0.2447528	0.6584392	0.0968080	0.0000000
							0.1046666	0.1046666	0.1046666	0.6860001
							0.2936310	0.2936310	0.2936310	0.1191069
							0.1141973	0.1141973	0.4885725	0.2830329
							0.1141973	0.1141973	0.4885725	0.2830329

$n$	$N_n$	$n_1$	$n_4$	$n_6$	$n_{12}$	$n_{24}$	$b_1$	$b_2$	$b_3$	$b_4$
8	165	1	1	1	2	0	1.0000000	0.0000000	0.0000000	0.0000000
							0.9533797	0.0466203	0.0000000	0.0000000
							0.8375919	0.1624081	0.0000000	0.0000000
							0.6801403	0.3198597	0.0000000	0.0000000
							0.5000000	0.5000000	0.0000000	0.0000000
							0.0627331	0.0627331	0.8745338	0.0000000
							0.2153606	0.2153606	0.5692789	0.0000000
							0.3891297	0.3891297	0.2217406	0.0000000
							0.3657423	0.5524728	0.0817849	0.0000000
							0.1942206	0.7294168	0.0763626	0.0000000
							0.2500000	0.2500000	0.2500000	0.2500000
							0.0834511	0.0834511	0.0834511	0.7496468
							0.4060462	0.4060462	0.0939538	0.0939538
							0.0927818	0.0927818	0.5844475	0.2299889
							0.2432058	0.2432058	0.4157364	0.0978521
9	220	0	2	0	4	0	1.0000000	0.0000000	0.0000000	0.0000000
							0.9626819	0.0373181	0.0000000	0.0000000
							0.8672666	0.1327334	0.0000000	0.0000000
							0.7361751	0.2638249	0.0000000	0.0000000
							0.5815151	0.4184849	0.0000000	0.0000000
							0.3333333	0.3333333	0.3333333	0.0000000
							0.0493729	0.0493729	0.9012542	0.0000000
							0.4658361	0.4658361	0.0683277	0.0000000
							0.1769439	0.1769439	0.6461122	0.0000000
							0.3020146	0.6309227	0.0670627	0.0000000
							0.1575680	0.7808733	0.0615587	0.0000000
							0.3261032	0.4887991	0.1850977	0.0000000
							0.0682526	0.0682526	0.0682526	0.7952422
							0.2123036	0.2123036	0.2123036	0.3630891
							0.0774288	0.0774288	0.6543249	0.1908175
							0.0781245	0.0781245	0.5016150	0.3421360
							0.2049128	0.2049128	0.5083123	0.0818620
							0.3552639	0.3552639	0.2073089	0.0821632

## References

- [1] Bernardi, C. and Maday, Y. (1992), *Approximations Spectrales de Problèmes aux Limites Elliptiques*. Springer-Verlag, Paris.
- [2] Bos, L.P. (1983), Bounding the Lebesgue Functions for Lagrange Interpolation in a Simplex. *J. Approx. Theo.* 38, 43-59.
- [3] Chen, Q. and Babuška, I. (1995), Approximate Optimal Points for Polynomial Intepolation in an Interval and in a Triangle.
- [4] Chung, K and Yao, T. (1977), On Lattices Admitting Unique Lagrange Interpolations, *SIAM J. Numer. Anal.* 14, 735-743.
- [5] Hughes, T.J.R. (1987), *The Finite Element Method: Linear Static and Dynamic Finite Element Analysis*. Prentice-Hall, Englewood Cliffs, NJ.
- [6] Sendov, Bl. and Andreev, A. (1994), Approximation and Interpolation Theory. in: Ciarlet, P.G. and Lions, J.L., eds., *Handbook of Numerical Analysis*, Vol. III, North-Holland, Amsterdam, 223-462.
- [7] Szabó, B.A. and Babuška, I. (1991), *Finite Element Analysis*, Wiley, New York.

**The Laboratory for Numerical Analysis** is an integral part of the Institute for Physical Science and Technology of the University of Maryland, under the general administration of the Director, Institute for Physical Science and Technology. It has the following goals:

To conduct research in the mathematical theory and computational implementation of numerical analysis and related topics, with emphasis on the numerical treatment of linear and nonlinear differential equations and problems in linear and nonlinear algebra.

To help bridge gaps between computational directions in engineering, physics, etc., and those in the mathematical community.

To provide a limited consulting service in all areas of numerical mathematics to the University as a whole, and also to government agencies and industries in the State of Maryland and the Washington Metropolitan area.

To assist with the education of numerical analysts, especially at the postdoctoral level, in conjunction with the Interdisciplinary Applied Mathematics Program and the programs of the Mathematics and Computer Science Departments. This includes active collaboration with government agencies such as the National Institute of Standards and Technology.

To be an international center of study and research for foreign students in numerical mathematics who are supported by foreign governments or exchange agencies (Fulbright, etc.).